

**NASA Contractor Report 181697**

**ICASE REPORT NO. 88-45**

# ICASE

THE STRONG NONLINEAR INTERACTION OF TOLLMIEEN-SCHLICHTING  
WAVES AND TAYLOR-GORTLER VORTICES IN CURVED CHANNEL FLOW

(NASA-CR-181697) THE STRONG NONLINEAR INTERACTION OF TOLLMIEEN-SCHLICHTING WAVES AND TAYLOR-GORTLER VORTICES IN CURVED CHANNEL FLOW Final Report (NASA) 35 p N88-29716  
CSCL 01B G3/01 0161713 Unclas

J. Bennett

P. Hall

F. T. Smith

Contract Nos. NAS1-18107 and NAS1-18605  
August 1988

INSTITUTE FOR COMPUTER APPLICATIONS IN SCIENCE AND ENGINEERING  
NASA Langley Research Center, Hampton, Virginia 23665

Operated by the Universities Space Research Association



National Aeronautics and  
Space Administration

Langley Research Center  
Hampton, Virginia 23665

# THE STRONG NONLINEAR INTERACTION OF TOLLMIEN-SCHLICHTING WAVES AND TAYLOR-GÖRTLER VORTICES IN CURVED CHANNEL FLOW

J. Bennett and P. Hall  
Mathematics Department  
University of Exeter  
Exeter, EX4 4QE, U.K.

F. T. Smith  
University College London  
Gower Street, London, WC1E 6BT, U.K.

## ABSTRACT

Viscous fluid flows with curved streamlines can support both centrifugal and viscous travelling wave instabilities. Here the interaction of these instabilities in the context of the fully developed flow in a curved channel is discussed. The viscous (Tollmien-Schlichting) instability is described asymptotically at high Reynolds numbers and it is found that it can induce a Taylor-Görtler flow even at extremely small amplitudes. In this interaction, the Tollmien-Schlichting wave can drive a vortex state with wavelength either comparable with the channel width or the wavelength of lower branch viscous modes. The nonlinear equations which describe these interactions are solved for nonlinear equilibrium states.

---

This work was supported by the National Aeronautics and Space Administration under NASA Contract Nos. NAS1-18107 and NAS1-18605 while the second and third authors were in residence at the Institute for Computer Applications in Science and Engineering (ICASE), NASA Langley Research Center, Hampton, VA 23665.

## 1. Introduction.

There are many fluid flows of practical importance where destabilizing centrifugal and viscous instability mechanisms are both present. Thus, for example, the flow in a curved rectangular duct or the flow over parts of a laminar flow wing (see Harvey and Pride (1982)) can support both Taylor-Görtler vortices and Tollmien-Schlichting waves. Indeed the latter flow can in turn support Rayleigh instabilities associated with the highly inflectional velocity profiles in some directions. In these flows the possibility exists that the nonlinear interaction of the different instability mechanisms might produce premature transition to turbulence. In that regard a potentially significant result from the recent work of Hall and Smith (1988a) is that interacting oblique Tollmien-Schlichting waves can generate longitudinal vortices essentially identical to Taylor-Görtler vortices even in the absence of wall curvature.

One of the first calculations on the interaction of Tollmien-Schlichting (TS) waves and Görtler vortices was given by Nayfeh (1981). Nayfeh discussed the effect of a Görtler vortex of given size on the growth of oblique TS waves. The amplitude of the vortex was assigned arbitrarily however and an eigenfunction shape found by solving the parallel-flow-approximation Görtler-vortex equations. It is now known (Hall 1982a, 1983) that, in the wavenumber regimes considered by Nayfeh, these equations have solutions of no relevance to spatially growing vortices. Furthermore, the amplitude of the vortex cannot be assigned arbitrarily; it must of course be determined by either a numerical or analytical solution of the Navier-Stokes equation as in Hall (1982b, 1988). Having made these approximations Nayfeh found that the vortices could have a massive effect on the growth of TS waves. Later Malik (1988) showed that Nayfeh's numerical calculations were incorrect and that his conclusions were in error.

More recently Bennett and Hall (1988) examined the effect of finite amplitude Görtler vortices in fully developed flows on the growth of lower branch TS waves. Here the nonlinear vortex state was found by solving the Navier-Stokes equations and a linear stability analysis of the solutions was given. It was shown that even small amplitude vortices can have a significant effect on TS growth rates. The vortex flow found by Bennett and Hall had a wavelength comparable to the depth of the channel in which the flow occurred. The asymptotic structure of TS waves appropriate to the lower branch of the neutral curve corresponds in general to TS waves with a small spanwise wavenumber and Bennett and Hall showed how this structure could be modified to allow for a faster spanwise dependence induced by the vortices .

The first nonlinear description of the interaction of vortices and TS waves was given by Hall and Smith (1988a). Here it was shown that long wavelength vortices and oblique

TS waves undergo a resonant triad interaction at small amplitudes. The interaction is, in the first instance, governed by ordinary differential triad amplitude equations which possess a finite time singularity. At higher amplitude the interaction is controlled by a coupled partial differential-ordinary integro-differential equation system. The solution of the system was found to depend crucially on the orientation of the TS waves to the vortices. The most dangerous type of interaction concerns TS waves propagating in a direction making an angle of more than  $41.6^\circ$  to the vortices since the resulting interaction produces a finite time singularity.

In this paper we shall investigate the stronger type of interaction which occurs after this singularity develops. The size of the TS waves is fixed by the condition that the downstream velocity component is perturbed at zeroth order by the vortex induced by the interacting TS waves. Thus this interaction leads to an  $O(1)$  change from the basic state that exists in the absence of vortices. If the size of the TS wave is decreased then a weakly nonlinear bifurcation governed by a cubic order amplitude equation typical of those found when using the Stuart-Watson method is retrieved. Surprisingly the stronger type of interaction can occur at extremely small TS amplitudes and both with and without wall curvature being present. The fact that such a relatively small TS wave can have such a large effect on the basic state is due to the large initial forces associated with the small streamwise length scale of the waves. The nonlinear equilibrium states appropriate to this interaction must be found numerically. A similar extension of the Bennett and Hall work into the nonlinear regime is also discussed for vortex flows of cross-stream wavelength comparable to the channel depth. We note here that other nonlinear states are accessible by routes other than that described in this paper; see, for example, Hall and Smith (1988b) where the vortex-TS interaction for external flows is discussed. The question of whether the present route described is physically the most relevant or whether one of the 'by-pass' routes dominates in an experiment cannot yet be answered.

In addition, at small vortex wavenumbers the three-dimensional breakdown of a 2D-TS wave can be described by an analysis of our interaction equations. Thus we determine the size of 2D-TS waves which are neutrally stable to oblique TS waves. A key feature of this secondary instability process is the longitudinal vortex system induced by the interacting oblique waves. Furthermore the interaction can occur in a straight channel, thus yielding a mechanism for the breakdown of 2D-TS waves in parallel flows.

The procedure adopted in the rest of this paper is as follows: in §2 the nonlinear equations governing fully nonlinear vortex flows in curved channels are described; in §3 these equations are solved in the presence of vortices of  $O(1)$  cross-stream wavenumber and the possible equilibrium states are described; in §4 the corresponding calculation for

small wavenumber vortices is described; and in §5 the instability of 2D-TS waves in a straight channel is discussed through the interaction equations derived in §4. Finally in §6 we discuss our results and draw some conclusions, mainly that nonlinear effects lead to a supercritical bifurcation to a mixed vortex-TS state.

## 2. Formulation of the disturbance equations for curved channel flows.

Consider the flow of a viscous incompressible fluid in a curved channel with walls defined by  $r^* = a, a + d$  with respect to cylindrical polar co-ordinates  $(r^*, \theta^*, z^*)$ . It is assumed that the curvature parameter,  $\delta$ , defined by

$$\delta = \frac{d}{a}, \quad (2.1)$$

is small. The flow is driven by the streamwise pressure gradient

$$\frac{1}{\rho} \frac{\partial p^*}{\partial \theta^*} = \frac{-12\nu V_m r^*}{d^2}, \quad (2.2)$$

where  $\rho$  and  $\nu$  are the density and kinematic viscosity of the fluid respectively, whilst  $V_m$  is a typical streamwise flow speed. The pressure gradient (2.2) drives a velocity field  $v_m$  in the streamwise direction and

$$v_m = V_m U(y), \quad (2.3)$$

where for small values of  $\delta$

$$U(y) = U_0(y) + O(\delta), \quad (2.4)$$

with

$$U_0(y) = 6y(1-y), \quad y = \frac{r^* - a}{d}. \quad (2.5a, b)$$

The dimensionless variables  $x, z$ , and  $t$  are then defined by

$$x = \frac{a\theta^*}{a}, \quad z = \frac{z^*}{d}, \quad t = \frac{t^*\nu}{d^2}. \quad (2.6a, b, c)$$

where  $t^*$  denotes the (dimensional) time. If  $(u, v, w)$  is the velocity field scaled on  $V_m$  with respect to  $(x, y, z)$  and the pressure is scaled on  $\rho V_m^2$  then the Navier-Stokes equations can be written in the form

$$\begin{aligned} \frac{1}{\mathcal{F}} \frac{\partial u}{\partial x} + \frac{\delta u}{\mathcal{F}} + \frac{\partial v}{\partial y} + \frac{\partial w}{\partial t} &= 0, \\ \frac{1}{Re} \{ \nabla^2 - \partial_t - \frac{\delta^2}{\mathcal{F}^2} \} u + \frac{2\delta}{Re\mathcal{F}^2} \frac{\partial v}{\partial x} - \frac{1}{\mathcal{F}} \frac{\partial p}{\partial x} &= Nu + \frac{\delta uv}{\mathcal{F}}, \\ \frac{1}{Re} \{ \nabla^2 - \partial_t - \frac{\delta^2}{\mathcal{F}^2} \} v - \frac{2\delta}{Re\mathcal{F}^2} \frac{\partial u}{\partial x} - \frac{\partial p}{\partial y} &= Nv - \frac{\delta u^2}{\mathcal{F}}, \\ \frac{1}{Re} \{ \nabla^2 - \partial_t \} w - \frac{\partial p}{\partial z} &= Nw, \end{aligned} \quad (2.7a, b, c, d)$$

where the Reynolds number  $Re$  is defined by

$$Re = V_m \frac{d}{\nu} \quad (2.8)$$

and

$$\mathcal{F} = 1 + \delta y,$$

$$\nabla^2 \equiv \frac{1}{\mathcal{F}^2} \partial_x^2 + \partial_y^2 + \frac{\delta}{\mathcal{F}} \partial_y + \partial_z^2, \quad (2.8b, c, d)$$

$$N = (u/\mathcal{F}) \partial_x + v \partial_y + w \partial_z.$$

The Taylor number  $T$  is then defined by

$$T = 4Re^2 \delta, \quad (2.9)$$

and it is known from the work of Dean (1928) that instability in the form of Taylor vortices occurs first for  $O(1)$  spanwise wavelengths with  $T$  also an  $O(1)$  quantity. In view of the smallness of  $\delta$  it follows from (2.9) that Taylor vortices first occur at high values of the Reynolds number. The flow is therefore susceptible to lower branch Tollmien-Schlichting instabilities with wavenumbers of order  $Re^{-\frac{1}{7}}$  in the azimuthal direction. It is the nonlinear interaction of these two modes of instability which will be discussed in the next section.

However, because a primary aim of the present calculation is to shed light on the related external flow problem where long wavelength Görtler vortices are important in the initial development of the flow, we consider a closely related interaction problem arising from the weakly nonlinear theory of Hall and Smith (1988a). Here the Taylor vortices occur with axial wavenumbers  $O(Re^{-\frac{1}{7}})$  which are comparable with the wavenumbers of lower branch Tollmien-Schlichting waves. These long wavelength vortices occur at relatively high values of the Taylor number  $T \sim O(Re^{\frac{2}{7}})$  which requires that  $Re \sim O(\delta^{-\frac{7}{12}})$  whereas the previous interaction has  $Re \sim O(\delta^{-\frac{1}{2}})$ . Thus, for a given value of  $Re$ , the former type of interaction occurs first when  $\delta$  is increased from zero. Alternatively, for a given channel with  $\delta$  fixed, the former interaction occurs at the lowest values of  $Re$ . However there is some indication from the work of Hall and Smith (1988a,b) that the second interaction can occur even at zero values of  $T$ , and so it is important to understand its structure.

Finally in this section we write down the form of the solutions of (2.7) appropriate to fully nonlinear Taylor vortices. Firstly, for  $O(1)$  vortex wavelengths the pressure  $p$  and velocity field  $(u, v, w)$  can be written as

$$p = \left[ -\frac{12x}{Re} + \frac{\tilde{p}}{2Re^2} \right] [1 + O(Re^{-1})] \quad (2.10a, b)$$

$$(u, v, w) = \left[ U_0 + \tilde{u}, \frac{\tilde{v}}{2Re}, \frac{\tilde{w}}{2Re} \right] [1 + O(Re^{-1})].$$

Here  $\tilde{u}, \tilde{v}, \tilde{w}, \tilde{p}$  are functions of  $t, y, z$  and satisfy

$$\begin{aligned}\frac{\partial \tilde{v}}{\partial y} + \frac{\partial \tilde{w}}{\partial z} &= 0 \\ [\nabla_2^2 - \partial_t] \tilde{u} - \frac{1}{2} \tilde{v} U_0' &= \frac{1}{2} (\tilde{v} \tilde{u}_y + \tilde{w} \tilde{v}_z) \\ [\nabla_2^2 - \partial_t] \tilde{v} - \frac{\partial \tilde{p}}{\partial y} + \frac{T}{2} \tilde{u} U_0 &= \frac{1}{2} (\tilde{v} \tilde{v}_y + \tilde{w} \tilde{v}_z) \\ [\nabla_2^2 - \partial_t] \tilde{w} - \frac{\partial \tilde{p}}{\partial z} &= \frac{1}{2} (\tilde{v} \tilde{w}_y + \tilde{w} \tilde{w}_z) \\ \tilde{u} = \tilde{v} = \tilde{w} &= 0, \quad y = 0, 1.\end{aligned}\tag{2.11a, b, c, d, e}$$

with  $\nabla_2^2 = \partial_y^2 + \partial_z^2$ .

The solution of the linearized form of the above equations shows that Taylor vortices grow exponentially in time for  $T > T_c \sim 5162$ . and weakly nonlinear theory (Seminara (1975)) can then be used to show that the most unstable mode is stabilized by nonlinear effects. The neutral curve for the linear stability problem is shown in Figure 1.

In the fully nonlinear regime possible stable finite amplitude solutions of (2.11) can be found by stepping the equations forward in time until the flow equilibrates. Bennett and Hall (1988) investigated the instability of these vortex flows to small Tollmien-Schlichting waves. Here in contrast we allow the Tollmien-Schlichting waves to be sufficiently large that they have an  $O(1)$  effect on the vortex flows.

Second, if the vortex wavelength is  $O(Re^{-\frac{1}{7}})$  we define

$$Z = Re^{-\frac{1}{7}} z, \quad \delta = \frac{1}{4} \tilde{T} Re^{-\frac{12}{7}},\tag{2.12a, b}$$

and (2.10) is modified to give

$$\begin{aligned}p &= \left[ \frac{-12x}{Re} + \frac{\tilde{p}}{2Re^{\frac{12}{7}}} \right] [1 + O(Re^{-\frac{5}{7}})], \\ [u, v, w] &= [U_0 + \tilde{u}, \frac{\tilde{v}}{2Re}, \frac{\tilde{w}}{2Re^{\frac{5}{7}}}] [1 + O(Re^{-\frac{5}{7}})].\end{aligned}\tag{2.13a, b}$$

The functions  $\tilde{u}$ ,  $\tilde{v}$ ,  $\tilde{w}$  and  $\tilde{p}$  are now functions of  $y$ ,  $Z$ , and  $t$  and satisfy

$$\begin{aligned}\frac{\partial \tilde{v}}{\partial y} + \frac{\partial \tilde{w}}{\partial Z} &= 0, \\ \{\partial_y^2 - \partial_t\} \tilde{u} &= \frac{1}{2} \tilde{v} U'_0 + \frac{1}{2} (\tilde{v} \tilde{u}_y + \tilde{w} \tilde{u}_Z), \\ \frac{1}{2} \tilde{T} \tilde{u} U_0 - \frac{\partial \tilde{p}}{\partial y} &= 0, \\ \{\partial_y^2 - \partial_t\} \tilde{w} - \frac{\partial \tilde{p}}{\partial Z} &= \frac{1}{2} \{\tilde{v} \tilde{w}_y + \tilde{w} \tilde{w}_Z\}, \\ \tilde{u} = \tilde{v} = \tilde{w} &= 0, \quad y = 0, 1.\end{aligned}\tag{2.14a, b, c, d, e}$$

The linearized form of this system has unstable disturbances for any nonzero value of  $\tilde{T}$ . Indeed the neutral value of  $\tilde{T}$  is proportional to  $k^{-2}$  if  $k$  is the axial wavenumber. This means that finite amplitude solutions of (2.14) cannot be found by integrating forward in time since the energy of the disturbance will rapidly cascade into the higher harmonics. We return to the long wavelength limit in §4, since we concentrate next on the nonlinear interaction problem for  $\frac{\partial}{\partial z} = 0(1)$ .

### 3. The nonlinear evolution equations for TS waves and Taylor vortices of $0(1)$ wavelength.

Since the only major difference between the present calculation and that of Bennett and Hall (1988) is that the TS wave now has an  $0(1)$  effect on the vortex only the essential details of the flow structure will be given. The size of the TS wave is fixed by the requirement that the nonlinear terms in the  $y$  and  $z$  momentum equations be comparable with those in (2.7c,d). This forcing is most important away from the viscous wall layers of thickness  $Re^{-\frac{2}{3}}$  in which the TS waves adjust to the no-slip condition at the wall. It is convenient at this stage to define the small parameter  $\epsilon$  by

$$\epsilon = Re^{-\frac{1}{3}}.$$

Suppose then that TS wave is proportional to

$$E = \exp[i(\alpha \epsilon x - \int^t \frac{\Omega(t) dt}{\epsilon^4})]$$

where the wavenumber  $\alpha$  and the slowly varying frequency  $\Omega$  are real. The frequency and amplitude of the wave must vary on the vortex time scale in order to allow for the situation when the vortex flow is evolving in time. If the vortex flow is in equilibrium then the frequency and amplitude of the TS wave are constant.



Away from the viscous layers near  $y = 0, 1$  the velocity field and pressure expand as

$$u = \left\{ \begin{pmatrix} U_0 + u_0 \\ \epsilon^7/2 \\ \epsilon^7 w_0/2 \end{pmatrix} + \left[ \begin{pmatrix} \epsilon^6 u_1 \\ \epsilon^7 v_1 \\ \epsilon^7 w_1 \end{pmatrix} E + \text{complex conjugate} \right] \left[ 1 + O(\epsilon^7) \right] \right\}. \quad (3.1a, b)$$

$$p = (-12x/Re + p_0/2Re^2 + [\epsilon^8 p_1 E + \text{complex conjugate}]) + O(\epsilon^7).$$

The above expansions are substituted into (2.7) and the zeroth order system for the  $x$  independent part of the flow satisfies the vortex equations (2.11) with  $(\tilde{u}, \tilde{v}, \tilde{w}, \tilde{p})$  replaced by  $(u_0, v_0, w_0, p_0)$  and with forcing terms  $F_1, F_2$  on the right hand sides of (2.11c,d). These forcing terms come from the interaction of the TS wave with itself and are given by

$$F_1 = 2[(-i\alpha u_1 v_1^* + v_1 v_{1y}^* + w_1 v_{1z}^*) + \text{complex conjugate}], \quad (3.2a, b)$$

$$F_2 = 2[(-i\alpha u_1 w_1^* + v_1 w_{1y}^* + w_1 w_{1z}^*) + \text{complex conjugate}].$$

(In these two equations, \* denotes complex conjugation.) The boundary conditions for  $(u_0, v_0, w_0, p_0)$  come from the no-slip condition

$$u_0 = v_0 = w_0 = 0 \text{ on } y = 0 \text{ and } 1.$$

together with a periodicity condition in  $z$ . This follows from the fact that the forcing terms become negligible in the wall layers.

Now consider the zeroth order approximation to the TS wave equations in the core. We find that they are identical to the corresponding equations in Bennett and Hall's linear analysis, so that writing  $\bar{U} = u_0 + U_0$  we have

$$i\alpha u_1 + v_{1y} + w_{1z} = 0, \quad i\alpha \bar{U} u_1 + v_1 \bar{U}_y + w_1 \bar{U}_z = 0 \quad (3.3a, b, c, d)$$

$$i\alpha \bar{U} v_1 = -p_{1y}, \quad i\alpha \bar{U} w_1 = -p_{1z},$$

together with the tangential slipping condition at the walls

$$v_1 = 0, \quad y = 0, 1.$$

Bennett and Hall simplified these equations by introducing a normalized pressure term  $\phi$ :

$$p_1 = p_{10}(t) + (i\alpha)^2 A(t) \phi(y, z)$$

$$u_1 = A \nabla^2 \phi / 2\bar{U}, \quad v_1 = -i\alpha A \phi_y / \bar{U}, \quad w_1 = -i\alpha A \phi_z / \bar{U},$$

where  $p_{10}(t)$  and the amplitude  $A(t)$  need to be determined and  $\phi$  satisfies the core equation

$$\left( \frac{\phi_y}{\bar{U}^2} \right)_y + \left( \frac{\phi_z}{\bar{U}^2} \right)_z = 0. \quad \phi \text{ periodic in } z. \quad (3.4)$$

$$\phi = 0 \text{ on } y = 0. \quad \phi = 1 \text{ on } y = 1.$$

A related coreflow problem arises in Smith (1977). We can now eliminate  $u_1, v_1$  and  $w_1$  from the forcing terms in the vortex equations to give

$$\begin{aligned} F_1 &= 2\alpha^2 |A|^2 \left\{ \left[ \frac{(\phi_y)^2 + (\phi_z)^2}{\bar{U}^4} \right] (\bar{U}^2)_y + \left[ \frac{(\phi_y)^2 + (\phi_z)^2}{\bar{U}^2} \right]_y \right\}, \\ F_2 &= 2\alpha^2 |A|^2 \left\{ \left[ \frac{(\phi_y)^2 + (\phi_z)^2}{\bar{U}^4} \right] (\bar{U}^2)_z + \left[ \frac{(\phi_y)^2 + (\phi_z)^2}{\bar{U}^2 (\phi_z)^2} \right]_z \right\}. \end{aligned} \quad (3.5a, b)$$

We note that the last terms in these equations can, if necessary, be absorbed into the pressure derivatives in the  $y, z$  vortex equations. In the two-dimensional case  $\bar{U} = \bar{U}(y)$ ,  $\phi = \phi(y)$  and we see that  $F_2 \equiv 0$ , and (using equation (3.5)) we note that  $F_1$  can be completely absorbed into  $p_{0y}$ . Thus for flows that are strictly two-dimensional nonlinearity occurs only at larger values of  $\Delta$ , as in Hall and Smith (1982).

Because the leading order wave equations are inviscid it is necessary to consider the flow within two viscous wall layers at  $y = 0, 1$ . The flow in these regions is exactly the same as in linear theory since the TS waves are sufficiently small. Thus

$$y = \epsilon^2 Y$$

in the lower viscous layer and

$$\begin{aligned} \underline{u} &= \{(\epsilon^2 \lambda_0(z, t) Y, -\epsilon^{11} \mu_{0z}(z, t) Y^2/2, \epsilon^9 \mu_0(z, t) T) \\ &+ [(\epsilon^6 U, \epsilon^9 V, \epsilon^7 W) E + \text{complex conjugate}] + \dots\}, \\ p^* &= \rho U_m^2 \{-12\epsilon^7 x - \epsilon^{16} \mu_{0z} Y \\ &+ [(\epsilon^8 P + \epsilon^{10} Q E + \text{complex conjugate})] + \dots\}. \end{aligned} \quad (3.6a, b)$$

where  $\lambda_0$  can be determined from  $\lambda_0(z, t) = \bar{U}_{|y=0}$ . The equations to determine  $U, V, W, P$  are

$$i\alpha U + V_y + W_z = 0.$$

$$i(-\Omega + \alpha \lambda_0 Y) U + \lambda_0 V + \lambda_{0z} Y W = -i\alpha P + U_{yy}, \quad (3.7)$$

$$P_y = Q_y = P_z = 0, i(-\Omega + \alpha \lambda_0 Y) W = -Q_z + W_{yy},$$

with boundary conditions  $U = V = W = 0$  on  $Y = 0$ , together with matching conditions with the core :

$$U \rightarrow A(t) \phi_{yyy|y=0}/(2\lambda_0), W \rightarrow 0 \quad \text{and} \quad P \rightarrow p_{10} \quad \text{as} \quad Y \rightarrow \infty,$$

and a periodicity condition. An important feature of these equations is the presence of a second order pressure term  $Q$ . We need to include this term because the  $z$  variation of the

flow is relatively fast compared to the  $x$  variation of the waves,  $O(1)$  as opposed to  $O(\epsilon)$ , so the  $z$ -derivatives of this smaller pressure will come in at higher order.

These boundary-layer equations were solved by Bennett and Hall (1988) in terms of Airy functions  $Ai(\xi)$  to give an eigen- or dispersion-relation. The equations governing the wall layer flow are then matched to the coreflow solution to give the following first order differential equation in  $Q_z$ ,

$$Q_{zz} + \frac{\lambda_{0z}}{\lambda_0} \psi Q_z = \alpha^2 p_{10} - \frac{1}{2} \alpha^2 A \phi_{vvv|v=0} \frac{Ai'(\xi)}{(i\alpha\lambda_0)^{\frac{1}{3}} K(\xi)}, \quad (3.8a)$$

where

$$\psi(\xi) = -\left(\frac{3}{2} + \frac{\xi}{2Ai(\xi)}[\xi K(\xi) + Ai'(\xi)]\right), \quad (3.8b)$$

$$\xi = -\frac{\Omega e^{i\pi/6}}{(\alpha\lambda_0)^{\frac{2}{3}}} \text{ and } K(\xi) = \int_{\xi}^{\infty} Ai(s) ds. \quad (3.8c, d)$$

A similar equation involving  $\lambda_1(z, t) = \bar{U}_{v|v=1}$  arises from the boundary-layer at  $y = 1$ . By applying the periodicity condition in  $z$ , combining the two results to eliminate  $p_{10}$  and  $A$  we can determine the eigenrelation. This is somewhat unweildy to write down but can best be calculated from the following:

$$\alpha^2 = \frac{1}{2} \sum_{n=0}^{n=1} \frac{f_n(2\pi/k)}{g_n(2\pi/k)}, \quad (3.9a)$$

where  $2\pi/k$  is the period of the vortices and  $f_n(z)$  and  $g_n(z)$  satisfy the following first order differential equations,

$$f_{nz} + \frac{\lambda_{nz}}{\lambda_n} \psi(\xi_n) f_n = \phi_{vvv|v=n} \frac{Ai'(\xi_n)}{(i\alpha\lambda_n)^{\frac{1}{3}} K(\xi_n)}. \quad (3.9b, c)$$

$$g_{nz} + \frac{\lambda_{nz}}{\lambda_n} \psi(\xi_n) g_n = 1,$$

with boundary conditions

$$f_n(0) = 0, g_n(0) = 0, \quad (3.9d)$$

with  $\psi$  and  $K$  defined above and  $\xi_n$  given by

$$\xi_n = -\frac{\Omega e^{i\pi/6}}{(\alpha\lambda_n)^{\frac{2}{3}}}, \quad n = 0, 1. \quad (3.9e)$$

In order to facilitate the numerical solution of the vortex equations it is convenient to

eliminate the pressure from the  $y$  and  $z$  momentum to give

$$\begin{aligned} \{\nabla^2 - \partial_t\}u_0 - \frac{1}{2}v_0U'_0 &= \mathcal{N}u_0, \\ \{\nabla^2 - \partial_t\}\nabla^2v_0 + TU_0u_{0zz} &= (\mathcal{N}v_0)_{zz} - (\mathcal{N}w_0)_{zz} - \frac{1}{2}T(u_0^2)_{zz} \\ &+ 2\alpha^2 |A|^2 \left\{ \frac{(\nabla\phi)_z^2(\bar{v})^2 - (\nabla\phi)_z^2(\bar{v})^2}{\bar{v}^4} \right\}_z, \\ u_0 = v_0 = w_0 &= 0, \quad y = 0, 1, \end{aligned} \tag{3.10}$$

where  $\mathcal{N} \equiv \frac{1}{2}\{v_0\partial_y + w_0\partial_z\}$  and  $\phi$  appearing on the right hand side of the equation for  $v_0$  satisfies the 'core' equation (3.4).

In the absence of a Tollmien-Schlichting wave it is well-known that, without any loss of generality, we can take  $u_0, v_0$  to be even functions of  $z$  and  $w_0$  is then an odd function of  $z$ . In fact (3.4), (3.10) and the wall equations show that the forced equation also admits a solution with  $u_0, v_0$  even in  $z$  with  $w_0$  odd in  $z$ . In this case  $Q$  and the corresponding upper wall pressure together with  $\phi$  are even functions of  $z$ . Thus at  $O(1)$  wavelengths the mixed vortex-TS state is found by solving (3.10) subject to the TS wall equations (3.9b,c).

### The numerical method.

When the Tollmien-Schlichting waves are small, as in the linear problem, the terms  $F_1$  and  $F_2$  become zero to leading order and the vortex and dispersion equations then decouple. Bennett and Hall solved equation (3.10) with  $F_1 \equiv F_2 \equiv 0$  using the method of Rogers and Beard (1969) for solving the Taylor vortex equations, by expanding the vortex velocities in Fourier sine and cosine series in  $z$  and by advancing in time to reach a steady solution. The nonlinear terms were calculated explicitly. Equation (3.4) was then solved for the wave pressure  $\phi$  by using finite differences in both  $y$  and  $z$  and iterating to a solution. Several thousand iterations were required, the exact number depending on the method used and the step lengths in  $y$  and  $z$ . Once the skin-frictions  $\lambda_0, \lambda_1$  and pressure derivatives  $\phi_{yyy}|_{y=0,1}$  had been determined  $\alpha$  and  $\Omega$  were found by solving the dispersion relation (3.9) using Runge-Kutta methods. The Airy functions were calculated using a combination of ordinary and asymptotic series, depending on the value of the argument  $\xi$ .

We could adapt this approach to solve our unsteady nonlinear problem in the following way. First we fix the T-S wave number  $\alpha$ , the initial vortex velocity  $\underline{u}_0(t_0)$  and wave pressure  $\phi(t_0)$ . We then guess an amplitude at the next time step  $|A(t_1)|$  and use it and  $\phi(t_0)$  to calculate the forcing terms  $F_1$  and  $F_2$ . The vortex equations (3.10) can then be solved for  $\underline{u}_0(t_1)$  while  $\phi(t_1)$  can be found from the pressure equation (3.4) using the new velocity. The eigenrelation (3.9) can then be solved for  $\Omega(t_1)$ . The guess  $A(t_1)$  can then be iterated on, until the imaginary part of  $\Omega$  is zero. This process could then be repeated

at the next time step  $t_2$ . We would then be able to observe the wave amplitude  $|A(t)|$  evolve in time for various wave-numbers  $\alpha$  away from the neutral point. The disadvantage with this method is that at each time step equation (3.4) needs to be solved accurately several times. The iterative method used by Bennett and Hall to solve (3.4) was very slow, taking up about half of the computing time for the entire linear calculation. We therefore feel a quicker scheme is needed for this part of the problem in order to carry out extensive numerical investigations using this approach.

It is possible however to calculate solutions to the steady nonlinear interaction problem less expensively. The method used is as follows. First we fix the term  $\alpha^2 |A|^2$  that occurs in equations (3.10). Starting with a guess for  $\underline{u}_0$  and  $\phi$  we march forward in time with the  $u_0, v_0$  equations. At each time step we update  $\phi$  from equation (3.4) by performing a few iterations. The forcing terms  $F_1$  and  $F_2$  can then be updated for the next time step. We continue this process until  $\underline{u}_0$  and  $\phi$  have settled down to steady solutions. The dispersion relation (3.9) can then be solved for real  $\alpha$  and  $\Omega$ . Once  $\alpha$  is known the steady amplitude  $|A|$  can be determined. Even if we were to solve the pressure equation (3.4) accurately at each time step, the velocities  $\underline{u}_0(t)$  that are produced before a steady solution emerges are not solutions of the physical problem. This is because we would find when we solved the eigenrelation (3.9) that in order to make  $\Omega$  real we would have to have different  $\alpha$ 's at each time step, whereas the theory requires  $\alpha$  to be independent of time. Only when the calculation has settled down to a steady solution do we have anything that makes sense physically. This means that the algorithm may well converge on unstable solutions as well as the stable ones, although of course we will not be able to tell which is which. Because  $\phi$  is only iterated upon a few times at each time step the overall time taken for this method is about equal to that for the linear calculation.

Since the TS wave part of the equations is still linear the calculation of  $\phi$  and the solution of the eigenrelation (3.9) are unchanged from Bennett and Hall. The algorithm used to step forward (3.4), (3.10) in time is virtually identical to that used by Bennett and Hall. The only difference is the presence in (3.10) of the forcing terms  $F_1$  and  $F_2$  which are periodic functions of  $z$  respectively. All the periodic functions of  $z$  can be expanded in the form

$$u_0 = \sum_{-\infty}^{\infty} u_{0n} e^{inkz} \quad (3.11)$$

where  $k$  is the vortex wavenumber. We recall that the equations from the interaction allow a solution with  $u_0, v_0$  and the TS wall pressure even in  $z$  and  $w_0, p_0$  odd in  $z$ . Thus we expect that there will be a solution of the nonlinear problem with  $u_{0n} = u_{0-n}$ , etc. However, in order to allow for the possibility of solutions without this symmetry we did not force this constraint. A consequence of this is that there is a mean (independent of

$x, z$ ) flow induced in the axial direction and an induced pressure gradient in that direction is needed to satisfy a mass flow constraint. A similar pressure function arises in the work of say DiPrima and Stuart (1975) or Hall (1984). Thus we allowed for such a function in our calculations. However, the only nonlinear solutions found had the above symmetries in  $z$  in which case there is no mean axial flow. It is possible that nonlinear solutions without these symmetries exist but they apparently do not bifurcate from the pure vortex flow. We postpone until §6 a discussion of the numerical results obtained following the scheme outlined above.

#### 4. The strongly nonlinear interaction between long wavelength Taylor vortices and Tollmien-Schlichting waves.

Here it is assumed that the channel curvature parameter  $\delta$  defined by (2.1) is sufficiently large to support Taylor vortices with cross-stream wavenumbers of order  $\epsilon = Re^{-\frac{1}{2}}$ . More precisely we consider the limits  $\delta \rightarrow 0, Re \rightarrow \infty$  with  $\tilde{T}$  defined by (2.12b) held fixed. In this case the basic circumferential flow can support vortices and oblique TS waves with comparable axial wavenumbers. The TS waves now have the structure first discussed by Smith (1978) in connection with the instability of the flow in an elliptical pipe. The overall size of the TS wave is again fixed by the condition that it should be sufficiently large in the core to drive the vortex flow at zeroth order and indeed alter the mean-flow profile. The streamwise and time dependence of the wave will again be expressed in terms of  $E = \exp i\{\alpha\epsilon x - \int^t \frac{\Omega(\tilde{t})}{\epsilon} dt\}$  in order to account for the possible evolution of the TS frequency as the vortex flow develops in time. The axial dependence of both modes is now of course entirely on the  $Z = \epsilon z$  scale. In this case the appropriate expansions of the velocity and pressure fields now become

$$\underline{u} = \left\{ \begin{pmatrix} U_0 + u_0 \\ \epsilon^7 v_0/2 \\ \epsilon^6 w_0/2 \end{pmatrix} + \cdots + \begin{pmatrix} \epsilon^5 u_1 \\ \epsilon^6 v_1 \\ \epsilon^7 w_1 \end{pmatrix} E + \text{complex conjugate} + \cdots \right\}, \quad (4.1a, b)$$

$$p = -12x\epsilon^7 + p_0 \frac{\epsilon^{12}}{2} + \cdots + \epsilon^7 p_1 E + \text{complex conjugate} + \cdots.$$

Here again the Taylor vortex functions  $u_0, v_0, w_0, p_0$  and the TS functions  $u_1, v_1, w_1, p_1$  must be matched with the appropriate expansions in the wall layers. In fact the Taylor vortex functions satisfy (2.14) but with forcing terms arising from the interaction of the Tollmien-Schlichting wave with itself. The zeroth order core equations for the TS wave yield the solution

$$\begin{aligned} u_1 &= A\bar{U}_y, v_1 = -i\alpha A\bar{U}, p_1 = -A\alpha^2 \int_{\frac{1}{2}}^y \bar{U}^2 dy, \\ w_1 &= \frac{\alpha^2 [A \int_{\frac{1}{2}}^y \bar{U}^2 dy]_x}{i\alpha \bar{U}}. \end{aligned} \quad (4.2a, b, c, d)$$

Here  $A$  is an amplitude function of  $Z = \epsilon z$  and the Taylor vortex time scale  $t$ . It should be noted that  $A$  cannot vary on a faster time scale more appropriate to a TS wave alone since the Taylor vortex cannot itself respond on such a scale induced by the forcing terms. With the form of the TS wave determined in the core the forcing terms in the vortex equations in the core can be expressed in terms of  $A$  and  $\bar{U}$ . If the vortex pressure is eliminated we obtain

$$\begin{aligned}
\frac{\partial v_0}{\partial y} + \frac{\partial w_0}{\partial Z} &= 0, \\
\{\partial_y^2 - \partial_t\} u_0 &= \frac{1}{2} v_0 U_{0y} + \mathcal{M} u_0, \\
\{\partial_y^2 - \partial_t\} \partial_y^2 v_0 + \tilde{T} U_0 u_{0ZZ} &= -(\mathcal{M} w_0)_{ZZ} - \frac{1}{2} \tilde{T} (u_0^2)_{ZZ} + 2\alpha^2 \{ |A|_Z^2 (\bar{U}^2)_y \}_Z, \\
u_0 = v_0 = w_0 &= 0, \quad y = 0, 1
\end{aligned} \tag{4.3}$$

where

$$\mathcal{M} \equiv v_0 \partial_y + w_0 \partial_Z.$$

We have anticipated above that the vortex flow in the presence of a TS wave should still satisfy the no-slip condition at the wall. In fact the axial momentum equation for the vortex flow in the wall layers is forced by the TS waves. However the forcing function decays to zero like the inverse square of the distance from the wall so that the appropriate matching conditions for  $w_0$  are that it should vanish at  $y = 0, 1$ .

It remains for us to determine the structure of the TS wave and Taylor vortex in the viscous wall layers. The appropriate expansions in the wall layer at  $y = 0$  are

$$\begin{aligned}
\underline{u} &= \{ [\epsilon^2 \lambda_0(Z, t) Y, \epsilon^{11} V_0, \epsilon^8 W_0] \\
&\quad [(\epsilon^5 U, \epsilon^8 V, \epsilon^5 W) E + \text{complex conjugate}] + \dots \} \\
p &= -12\epsilon^7 x + \epsilon^{12} P_0 + \dots + (\epsilon^7 P E + \text{complex conjugate}) + \dots
\end{aligned} \tag{4.4a, b}$$

The axial momentum equation for the vortex is now the only equation forced by the TS wave. However, the equations for the TS wave depend only on the vortex flow through the shear  $\lambda_0$  so we do not solve for  $V_0, W_0$  and  $P_0$  here. It suffices to say that the vortex equations in the wall layer at  $y = 0$  can be solved such that the solution matches with the coreflow vortex solution. The zeroth order equations for the TS waves in the wall layer at

$y = 0$  are

$$\begin{aligned}
i\alpha U + V_Y + W_Z &= 0 \\
i(-\Omega + \alpha\lambda_0 Y)U + \lambda_0 V + \lambda_{0Z} YW &= -i\alpha P + U_{YY} \\
P_Y &= 0 \\
i(-\Omega + \alpha\lambda_0 Y)W &= -P_Z + W_{YY}
\end{aligned} \tag{4.5a, b, c, d}$$

with boundary conditions

$$U = V = W = 0, \quad Y = 0 \tag{4.5e}$$

and matching condition

$$U \rightarrow \lambda_0 A, \quad Y \rightarrow \infty. \tag{4.5f}$$

Following Smith (1979) we can solve (4.5d) to give

$$W = -\frac{P_Z \mathcal{L}(\xi)}{\Delta^{\frac{2}{3}}}, \tag{4.6}$$

where

$$\begin{aligned}
\mathcal{L} &= A_i(\xi) \int_{\xi_0}^{\xi} \frac{ds}{A_i^2(s)} \int_{\infty}^s A_i(s_1) ds_1, \\
\xi &= \Delta^{\frac{1}{3}} \left( Y - \frac{\Omega}{\alpha\lambda_0} \right), \quad \xi_0 = \frac{-i\frac{1}{2}\Omega}{\Delta^{\frac{1}{3}}}.
\end{aligned}$$

The  $x$  velocity component of the TS wave is then given by

$$\begin{aligned}
\Delta^{\frac{5}{3}} U_{\xi} &= -\lambda_0 P_{ZZ} \mathcal{L}'(\xi) + \frac{1}{3} \lambda_{0Z} P_Z \{ 3\mathcal{L}'(\xi) + \frac{1}{2} \mathcal{L}''''(\xi) \\
&\quad - 2A_i'(\xi) \frac{\xi_0 \mathcal{L}'(\xi_0)}{A_i(\xi_0)} \} + B A_i(\xi),
\end{aligned} \tag{4.7}$$

which matches with the coreflow solution if

$$B = \frac{\Delta^{\frac{5}{3}} \lambda_0 A}{K(\xi_0)} - \frac{1}{2} \lambda_{0Z} P_Z \xi_0 \frac{\mathcal{L}'(\xi_0)}{K(\xi_0)}, \tag{4.8}$$

with  $K(\xi_0)$  defined by (3.8d). A similar expansion procedure can be carried out in the upper layer, while the core equations for the TS wave show that  $P$  and  $Q$ , the pressures in the lower and upper layers respectively, are related by

$$P - Q = \alpha^2 A \int_0^1 \bar{U}^2 dy. \tag{4.9}$$

The TS pressure  $P$  can be written using (4.5b) as

$$i\alpha P = \Delta^{\frac{2}{3}} U_{\xi \xi}|_{\xi=\xi_0}$$



so that (4.7), (4.8) now give

$$P'' + \frac{\lambda'_0}{\lambda_0} \psi_0 P' = \frac{A'_i(\xi_0) \lambda_0^2 [Q - P]}{K_0 \alpha^2 J} + \alpha^2 P, \quad (4.10)$$

where  $\psi_0$  is defined by (3.8b) with  $\xi = \xi_0$  and

$$J = \int_0^1 \bar{U}^2 dy. \quad (4.11)$$

A similar calculation in the upper layer shows that

$$Q'' + \frac{\lambda'_1}{\lambda_1} \psi_1 Q' = \alpha^2 Q + \frac{A'_{i0}(\xi_1) \lambda_1^2 [P - Q]}{K_0 \alpha^2 J}. \quad (4.12)$$

Here  $\lambda_1 = -\frac{\partial \bar{U}}{\partial y} \big|_{y=1}$  and  $\xi_1, \psi_1$  are obtained from  $\xi_0, \psi_0$  by replacing  $\lambda_0$  by  $\lambda_1$  in the definition of these quantities. Thus the evolution equations (4.3), (4.9), (4.10), (4.12) determine the TS and vortex structure as the nonlinear interactions take place. The vortex velocity components  $u_0, v_0, w_0$ , the Tollmien-Schlichting amplitude  $A$  (or  $P, Q$ ) and frequency  $\Omega$  are all functions of the vortex time scale  $t$ . Again it should be noted that the vortex structure cannot tolerate an evolution on a faster time scale typical of TS wave growth.

Suppose then that, for a given  $\alpha, \tilde{T}$ , we are given  $u_0, v_0, w_0, P$ , and  $\Omega$ . If a guess is made for  $|A|_{max}$  at the next time step (4.3) can be advanced using some finite difference discretization in  $t$  and  $y$  together with a Fourier expansion in  $Z$ . The wall equations (4.10), (4.12) at the new time step constitute an eigenrelation  $\Omega = \Omega(\alpha)$  and this can be solved for the complex  $\Omega$  which satisfies this eigenrelation. It is however necessary in our expansion procedure for  $\Omega$  to be purely real so the calculation is repeated with a different  $|A|_{max}$  in order to make  $\Omega_i = 0$ . At each stage of this iteration procedure  $A(z)$  is written as

$$A = |A|_{max} \mathcal{A}(z)$$

and  $\mathcal{A}(z)$  is taken from the solution of the eigenrelation at the new time step. When the iteration has converged the scheme can be used to advance the solution again. If there are stable equilibrium states ( $\partial_t = 0$ ) then we would expect that after a sufficiently long time the vortex and TS wave would become steady. The above procedure will then follow the time dependent evolution of the vortex-TS interaction. Because this calculation must be carried out over a significant time interval and there are many values of  $\alpha, \tilde{T}$  to be investigated it was decided to concentrate on an alternative scheme which would locate any equilibrium states directly. In this scheme  $\partial_t$  is set equal to zero and values for  $\alpha, |A|_{max}, \tilde{T}$  are chosen and a guess is made for  $\mathcal{A}(z)$ . We then use a Newton iteration procedure on the steady vortex equations to find the corresponding equilibrium vortex flow.

The TS wall equations are then solved for the real  $\alpha$  and  $\Omega$  which satisfy the eigenrelation. The new amplitude function  $A(z)$  obtained from this calculation is then used in the vortex equations to find an 'updated' vortex flow. The wall equations are then used to 'update'  $\alpha, \Omega$  and the procedure continued until the vortex solution and  $(\alpha, \Omega)$  converge for the given value of  $\alpha, |A|_{max}$  and  $\tilde{T}$ .

It was found that the above iteration procedure converged and typically about 100 steps in  $(0, 1)$  were taken to integrate the vortex equations using a Runge-Kutta scheme. The wall equations were solved by expanding  $P, Q$  and the coefficients in (4.10), (4.12) in Fourier series in  $Z$ . It was found that about 8 Fourier terms were usually adequate to solve the eigenrelation to the accuracy needed to plot the figures given later in this paper.

## 5. Secondary instability of two-dimensional Tollmien-Schlichting waves described by the TS-vortex interaction equations.

Here we demonstrate how the interaction equations can be used to model the three-dimensional breakdown of two-dimensional TS waves in a straight channel. Suppose then that  $A(Z, t)$ , the wave amplitude in the core region, takes the form

$$A(Z, t) = A_0(t) + A_1 e^{i\omega t} \cos \beta Z \quad (5.1)$$

where  $A_1$  is independent of time and is small compared with the 2D amplitude  $A_0$ . If  $A_1 = 0$  the TS forcing term in (4.3) vanishes identically so that  $A_0$  does not itself induce a longitudinal vortex field. This means that  $A_0(t)$  evolves as the solution of a cubic amplitude equation which must be found at higher order. In fact  $A_0(t)$  evolves on a longer time scale than  $t$  so that in (5.1)  $A_0$  can be treated as a constant. We shall determine the values of  $A_0$  at which the 3D perturbation in (5.1) is neutrally stable. The precise form of the perturbation in (5.1) corresponds to two equally inclined oblique TS waves of the same amplitude.

The vortex velocity field  $(u_0, v_0, w_0)$  induced by the TS wave (5.1) can be expressed as

$$(u_0, v_0, w_0) = [4\alpha^2 \beta^2 A_1 A_0 (u_{01}, v_{01}, w_{01}) \cos \beta Z e^{i\omega t} + \text{complex conjugate}] + O(A_1^2). \quad (5.2)$$

Here  $(u_{01}, v_{01}, w_{01})$  satisfy the problem

$$\begin{aligned} \frac{\partial v_{01}}{\partial y} - \beta w_{01} &= 0 \\ (\partial_y^2 - i\omega) u_{01} &= \frac{1}{2} v_{01} U'_0 \\ (\partial_y^2 - i\omega) \partial_y^2 v_{01} &= -U_0 U'_0 \\ u_{01} = v_{01} = v'_{01} &= 0, \quad y = 0, 1 \end{aligned} \quad (5.3)$$

which must be solved numerically (see below).

In the lower and upper wall layers  $P$  and  $Q$  expand as

$$P = P_0 + P_1 \cos \beta Z e^{i\omega t}$$

$$Q = Q_0 + Q_1 \cos \beta Z e^{i\omega t}$$

and from (4.9) it follows that

$$P_1 - Q_1 = \alpha^2 \left[ \frac{6}{5} A_1 + 8 A_1 A_0^2 \alpha^2 \beta^2 J_1 \right]$$

where

$$J_1 = \int_0^1 U_0 u_{01} dy.$$

In fact it can be shown that  $Q_1 = -P_1$ . If the wall equations are expanded in powers of  $A$ , then at zeroth order we obtain

$$\alpha^2 = \frac{5.6^{\frac{2}{3}} A'_{i0}}{K_0 \Delta^{\frac{1}{3}}} \quad (5.4)$$

which yields the usual neutral values of  $\alpha, \Omega$  when

$$\frac{A_{i0}}{\chi} \simeq 1.001, \xi = -2.298 i^{\frac{1}{3}}.$$

The condition (5.4) ensures that the two-dimensional TS wave is in neutral equilibrium. At next order we find that

$$1 + \frac{20}{3} \beta^2 \alpha^2 A_0^2 J_1 = \alpha^2 \Delta_0^2 \left[ \frac{-5}{3} J_1 + \frac{u'_{01}(0)}{2} \Phi \right] \quad (5.5)$$

where

$$\Phi = \frac{5}{18} - \frac{\xi A_{i0}}{9} \left( \frac{\xi_{01}}{A_{i0}} + \frac{1}{K_0} \right).$$

The integral  $J_1$  and the shear  $u_{01}(0)$  depend on  $\omega$  and (5.5) can be solved by taking real and imaginary parts to give

$$\beta^2 = 8b/d,$$

$$\alpha^2 A_0^2 = \frac{1}{8[a - \frac{b\xi}{a}]}$$

where

(5.6a, b, c, d)

$$-\frac{5}{6} J_1 + \frac{\Phi u'_{01}(0)}{2} = a + ib,$$

$$\frac{220}{3} J_1 = c + id.$$

A numerical investigation of (5.3) shows that (5.6a,b) give  $\beta^2 > 0, \alpha^2 A_0^2 > 0$  for  $\omega > 59.5$ . In fact the right hand side of (5.6b) is positive for  $\omega > 0$  while the right hand side of (5.6a)  $\rightarrow \infty$  when  $\beta\alpha^{-1} \rightarrow \infty$ , and so 3D waves oriented at all angles to the flow direction can be neutral. However, the value of  $A_0$  corresponding to  $\beta$  increases monotonically with  $\beta\alpha^{-1}$  so that 3D waves propagating almost parallel to the flow direction (i.e. with small  $\beta$ ) are, in a sense, the most dangerous. Thus (5.6a,b) determine the amplitude of the 2D TS wave which is neutrally stable to a pair of oblique waves with wavenumber  $\beta$ . It follows that at this amplitude there can be a secondary instability of the 2D TS wave to a pair of equally inclined oblique modes.

## 6. Results and Discussion.

We first discuss our results for the interaction problem of §3 which we recall concerns Görtler vortices of wavelength comparable with the channel width. In this case, finite amplitude vortex flows are possible only for  $T > T_c = 5162$  and the least stable axial wavenumber has  $k = 3.951$ . Since there are no unstable TS waves with streamwise wavenumbers  $O(Re^{-\frac{1}{2}})$  and spanwise wavenumbers  $O(1)$ , we restricted our calculations to the case  $T > T_c$  and spanwise wavenumber  $k = 3.951$ .

In Figures 1, 2, we show the dependence of  $\alpha |A|$  and  $\Omega$  on  $\alpha$  for equilibrium states corresponding to  $T = 11,000$ , and  $T = 22,000$  respectively. We see that at these Taylor numbers the neutral wavenumber and frequency both decrease monotonically as  $|\alpha A|$  increases. Results for  $T = 5500$  have the same properties and are shown in Table 1. From these results, we deduce that the bifurcation diagram for a fixed frequency disturbance is as sketched in Figure 3. We have not computed sufficient results to draw such a figure in detail because it would require an order-of-magnitude more computing time than that needed to generate Figures 1, 2. Linear theory shows that for each of the points of Figures 1, 2 the flow with infinitesimal  $|A|$  is unstable. We conclude that the interaction problem at  $O(1)$  vortex wavelengths leads to a supercritical bifurcation to a stable mixed TS-vortex state.

Now let us turn to our results for the interaction problem specified by (4.3), (4.9), (4.10) and (4.11) corresponding to a longer-scale spanwise dependence. In the absence of a finite amplitude Görtler vortex, we find that the neutral values of  $\alpha$  and  $\Omega$  corresponding to an oblique TS wave with  $A \sim \cos z$  or  $A \sim \sin z$  are

$$\alpha = 4.396 \quad , \quad \Omega = 20.456.$$

In the first instance we confine our remarks to the situation when  $\tilde{T} > 23711$  so that a finite amplitude vortex can exist in the absence of TS waves. In Figure 4 we show how

$\alpha, \Omega$  vary in the presence of a finite amplitude vortex at different values of the Taylor number. These values correspond to the linearized problem  $A \rightarrow 0$  and we see that both  $\alpha$  and  $\Omega$  increase monotonically with  $\tilde{T}$ . We note that a constant frequency TS wave will change from being stable to unstable when  $\tilde{T}$  is decreased through any point on the curve of Figure 4 for  $\tilde{T}$  greater than its linear neutral value  $\sim 23711$ .

Next, suppose that the Taylor number is held fixed and the neutral values of  $\alpha$  and  $\Omega$  are calculated for different values of  $|A|_{maz}$ . When  $|A|_{maz} \rightarrow 0$ , then the neutral values of  $\alpha$  and  $\Omega$  must tend back to the neutral values appropriate to the current value of the Taylor number. The results of such a calculation are shown in Figure 5 at two typical values of the Taylor number. We see that  $\alpha$  and  $\Omega$  increase monotonically when  $|A|_{maz}$  increases. We conclude that the bifurcation picture for any constant frequency TS wave will be as sketched in Figure 6. At first sight, it appears that this is a subcritical bifurcation to a mixed vortex-TS state. However, we found earlier that a constant frequency linear TS wave changes from being stable to unstable when  $\tilde{T}$  is increased, so that, if we associate the terms 'subcritical' and 'supercritical' with the linearly stable and unstable regions, we conclude that then a constant frequency disturbance undergoes a supercritical bifurcation to a finite amplitude state. A weakly nonlinear analysis of (4.3), (4.9), (4.10), (4.11) shows that when this bifurcation occurs the finite amplitude solution is stable and we do not expect that there will be any finite time singularities of the full time-dependent problem associated with the finite amplitude state.

Now let us consider the situation when  $\tilde{T} < 23711$  so that in the absence of a Tollmien-Schlichting wave there is no vortex activity. In order to see the differences which emerge in this case, we shall now indicate briefly how an amplitude equation typical of those obtained for weakly nonlinear stability problems using the Stuart-Watson method can be retrieved. Suppose then that the Taylor number  $\tilde{T}$  is held fixed and that the neutral values of  $\alpha$  and  $\Omega$  are given by

$$\alpha = \alpha_N, \quad \Omega = \Omega_N. \quad (6.1)$$

These are the neutral values appropriate to  $|A|_{maz} \rightarrow 0$ , so that if we now write

$$|A|_{maz} = \delta^{\frac{1}{2}}$$

where  $\delta$  is small and positive we anticipate that with  $\tilde{T}$  held fixed the appropriate expansions of  $\alpha$  and  $\Omega$  become

$$\begin{aligned} \alpha &= \alpha_N + \delta \tilde{\alpha} + \dots, \\ \Omega &= \Omega_N + \delta \tilde{\Omega} + \dots. \end{aligned} \quad (6.2a, b)$$

We see from (4.3) that  $u_0, v_0, w_0$  in (4.3) must be expanded as

$$u_0 = \delta \tilde{u} \cos 2z + \dots,$$

$$v_0 = \delta \tilde{v} \cos 2z + \dots,$$

$$w_0 = \delta \tilde{w} \sin 2z + \dots,$$

where  $\tilde{u}, \tilde{v}, \tilde{w}$  satisfy

$$\begin{aligned} \frac{d\tilde{v}}{dy} + 2\tilde{w} &= 0, \\ \frac{d^2\tilde{u}}{dy^2} &= \frac{1}{2}\tilde{v}U_{0y}, \\ \frac{d^4\tilde{v}}{dy^4} - 4\tilde{T}U_0 &= -4\alpha^2\tilde{T}U_{0y}^2, \end{aligned} \tag{6.3a, b}$$

$$\tilde{u} = \tilde{v} = \tilde{w} = 0 \quad , \quad y = 0, 1.$$

We note that the homogeneous form of (6.3) has a solution if  $4\tilde{T} \simeq 23711$ , so that  $\tilde{u}, \tilde{v}, \tilde{w}$  then become singular near  $\tilde{T} = \tilde{T}^+ \simeq 5978$ . In fact, these functions behave like  $\frac{1}{\tilde{T} - \tilde{T}^+}$  so that  $\hat{u}, \hat{v}, \hat{w}$  necessarily change sign at  $\tilde{T}^+$ . The wall equations (4.10) and (4.12) together with (4.11) are then found to have a solution only if  $\tilde{\alpha}, \tilde{\beta}$  satisfy an equation of the form

$$\tilde{\alpha}[c + id] + \tilde{\Omega}[e + if] + g + ih = 0. \tag{6.4}$$

Here  $c, d, e, f, g$  and  $h$  are real constants, the first four of which arise from the expansions of  $\alpha$  and  $\Omega$  in (4.10) - (4.12), whilst  $g$  and  $h$  arise from  $\hat{u}$ , the order  $\delta$  correction to the coreflow. By taking real and imaginary parts in (6.4) we can solve for  $\tilde{\Omega}$  and  $\tilde{\alpha}$ . Our numerical solution of the full problem for  $\tilde{T} > 23711$  suggests that there  $\Omega$  is always positive. However, because of the singularity of  $\tilde{u}$  at  $\tilde{T}^+$  it follows that  $\tilde{\Omega}$  must change sign at  $\tilde{T} = \tilde{T}^+$ .

It follows from the above that in the neighbourhood of  $\tilde{T}^+$  there is a change in the bifurcation structure of the mixed vortex-TS state; thus on one side of this Taylor number the bifurcation will be supercritical whilst on the other side it will be subcritical. In fact, this structure is discussed by Hall and Smith (1988a). In the present investigation the iteration scheme used to solve the interaction problem of §4 failed to converge for Taylor numbers less than 25000. Until the method failed to converge it was found that the bifurcation to the mixed state was always supercritical. This suggests that at sufficiently high Taylor numbers the time dependent interaction problem does not have a finite time singularity but that at lower Taylor numbers there could be such a singularity associated with the subcritical bifurcation.

## References

- [1] Bennet, J. and Hall, P. (1988), "On the secondary instability of Taylor-Görtler vortices to Tollmien-Schlichting waves in fully-developed flows," *J. Fluid Mech.*, Vol. 188, p. 445.
- [2] Dean, W. R. (1928), "Fluid motion in a curved channel," *Phil. Mag.*, Vol. 5, p. 673.
- [3] DiPrima, R. C. and Stuart, J. T. (1975), "The instability of the flow between eccentric cylinders," *J. Fluid Mech.*, Vol. 67, p. 85.
- [4] Hall, P. (1982), "Taylor-Görtler vortices in fully developed or boundary layer flows: Linear theory," *J. Fluid Mech.*, Vol. 124, p. 475.
- [5] Hall, P. (1982), "On the nonlinear evolution of Görtler vortices in growing boundary layers," *J. Inst. Maths. and Applications*, Vol. 29, p. 173.
- [6] Hall, P. (1984), "On the stability of the unsteady boundary layer on a cylinder oscillating transversely in a viscous fluid," *J. Fluid Mech.*, Vol. 146, p. 347.
- [7] Hall, P. (1988), "The growth of nonlinear Görtler vortices in growing boundary layers," *J. Fluid Mech.*, in press.
- [8] Hall, P. and Smith, F. T. (1982), "A suggested mechanism for nonlinear wall roughness effects on high Reynolds number flow stability," *Studies in Applied Mathematics*, Vol. 66, p. 241.
- [9] Hall, P. and Smith, F. T., (1988a), "The nonlinear interaction of Görtler vortices and Tollmien-Schlichting waves in curved channel flows," *Proc. Roy. Soc. Lond.*, Vol. A417, p. 244.
- [10] Hall, P. and Smith, F. T. (1988b), "Nonlinear Tollmien-Schlichting/vortex interaction in boundary layers," To appear as an ICASE report and submitted to *Journal of Fluid Mechanics*.
- [11] Harvey, W. D. and Pride, J. (1982), AIAA Paper No. 82-0567.
- [12] Malik, M. (1986), AIAA Paper No. 86-1129.
- [13] Nayfeh, A. H. (1981), "Effect of streamwise vortices on Tollmien-Schlichting waves," *J. Fluid Mech.*, Vol. 107, pp. 441-453.

- [14] Rogers, E. H. and Beard, D. W. (1969), "A numerical study of wide-gap Taylor vortices," J. Comp. Phys., Vol. 4, p. 1.
- [15] Seminara, G. (1975), "Instability of some unsteady viscous flows," Ph.D. Thesis, University of London, England.
- [16] Smith, F. T. (1979), "Instability of flow through pipes of general cross-section," Matematika, Vol. 26, pp. 187-210.



T	k	Amplitude	$\alpha$	$\Omega$
5500.	3.951	.1104	4.5289	20.7614
5500.	3.951	.1561	4.5284	20.7600
5500.	3.951	.1913	4.5279	20.7585
5500.	3.951	.2705	4.5266	20.7542




Fig. 1a. The dependence of  $|\alpha A|$  on  $\alpha$  for  $T = 11,000$ ,  $k = 3.951$ .




Fig. 1b. The dependence of  $|\alpha A|$  on  $\Omega$  for  $T = 11,000$ ,  $k = 3.951$ .




Fig. 2a. The dependence of  $|\alpha A|$  on  $\alpha$  for  $T = 22,000$ ,  $k = 3.951$ .




Fig. 2b. The dependence of  $|\alpha A|$  on  $\Omega$  for  $T = 22,000$ ,  $k = 3.951$ .




Fig. 3. A sketch of the bifurcation diagram for a constant frequency TS wave with  $k = 3.951$  for the vortex flow.




Fig. 4. The dependence of  $\alpha$  and  $\Omega$  on  $\tilde{T}$ .




Fig. 5a. The dependence of  $|A|_{\max}$  on  $\alpha$  for the case  $\tilde{T} = 28511$ .

Fig. 5b. The dependence of  $|A|_{\max}$  on  $\Omega$  for the case  $\tilde{T} = 28511$ .






Fig. 5c. The dependence of  $|A|_{\max}$  on  $\alpha$  for the case  $\tilde{T} = 31000$ .

Fig. 5d. The dependence of  $|A|_{\max}$  on  $\Omega$  for the case  $\tilde{T} = 31000$ .




Fig. 6. A sketch of the bifurcation diagram for a constant frequency TS wave.



## Report Documentation Page

1. Report No. NASA CR-181697 ICASE Report No. 88-45		2. Government Accession No.		3. Recipient's Catalog No.	
4. Title and Subtitle  THE STRONG NONLINEAR INTERACTION OF TOLLMIE- SCHLICHTING WAVES AND TAYLOR-GÖRTLER VORTICES IN CURVED CHANNEL FLOW				5. Report Date  August 1988	
				6. Performing Organization Code	
7. Author(s)  J. Bennett, P. Hall, and F. T. Smith				8. Performing Organization Report No.  88-45	
				10. Work Unit No.  505-90-21-01	
9. Performing Organization Name and Address  Institute for Computer Applications in Science and Engineering Mail Stop 132C, NASA Langley Research Center Hampton, VA 23665-5225				11. Contract or Grant No.  NAS1-18107, NAS1-18605	
				13. Type of Report and Period Covered  Contractor Report	
12. Sponsoring Agency Name and Address  National Aeronautics and Space Administration Langley Research Center Hampton, VA 23665-5225				14. Sponsoring Agency Code	
15. Supplementary Notes  Langley Technical Monitor: Submitted to Journal of Fluid Richard W. Barnwell Mechanics  Final Report					
16. Abstract  Viscous fluid flows with curved streamlines can support both centrifugal and viscous travelling wave instabilities. Here the interaction of these instabilities in the context of the fully developed flow in a curved channel is discussed. The viscous (Tollmien-Schlichting) instability is described asymptotically at high Reynolds numbers and it is found that it can induce a Taylor-Görtler flow even at extremely small amplitudes. In this interaction, the Tollmien-Schlichting wave can drive a vortex state with wavelength either comparable with the channel width or the wavelength of lower branch viscous modes. The nonlinear equations which describe these interactions are solved for nonlinear equilibrium states.					
17. Key Words (Suggested by Author(s))  longitudinal vortices, transition			18. Distribution Statement  01 - Aeronautics (General)  Unclassified - unlimited		
19. Security Classif. (of this report)  Unclassified	20. Security Classif. (of this page)  Unclassified		21. No. of pages  34	22. Price  A03	